

The articles below were used to develop [FACT's Nutritional Benefits of Pasture Raised Food handouts](#). Skip to: [beef cattle](#), [bison](#), [dairy cows](#), [laying hens](#), [meat birds](#), [pigs](#), and [sheep & goats](#).

Beef cattle

- Aaslyng, M. D., & Meinert, L. (2017). Meat flavour in pork and beef – From animal to meal. *Meat Science*, 132, 112–117. <https://doi.org/10.1016/j.meatsci.2017.04.012>
- Alfaia, C. M., Alves, S. P., Pestana, J. M., Madeira, M. S., Moreira, O., Santos-Silva, J., Bessa, R. J., Toldrá, F., & Prates, J. A. (2016). Distinct fatty acid composition of some edible by-products from bovines fed high or low silage diets. *Food Science and Technology International*, 23(3), 209–221. <https://doi.org/10.1177/1082013216674137>
- Avilés, C., Martínez, A., Domenech, V., & Peña, F. (2015). Effect of feeding system and breed on growth performance, and carcass and meat quality traits in two continental beef breeds. *Meat Science*, 107, 94–103. <https://doi.org/10.1016/j.meatsci.2015.04.016>
- Bermingham, E. N., Agnew, M., Gomes Reis, M., Taukiri, K., Jonker, A., Cameron-Smith, D., & Craigie, C. R. (2020). Assessment of atherogenic index, long-chain omega-3 fatty acid and phospholipid content of prime beef: a survey of commercially sourced New Zealand Wagyu and Angus beef cattle. *Animal Production Science*. Published. <https://doi.org/10.1071/an19427>
- Bjorklund, E., Heins, B., DiCostanzo, A., & Chester-Jones, H. (2014). Fatty acid profiles, meat quality, and sensory attributes of organic versus conventional dairy beef steers. *Journal of Dairy Science*, 97(3), 1828–1834. <https://doi.org/10.3168/jds.2013-6984>
- Blanco, M., Casasús, I., Ripoll, G., Albertí, P., Panea, B., & Joy, M. (2017). Is meat quality of forage-fed steers comparable to the meat quality of conventional beef from concentrate-fed bulls? *Journal of the Science of Food and Agriculture*, 97(14), 4943–4952. <https://doi.org/10.1002/jsfa.8371>
- Burnett, D. D., Legako, J. F., Phelps, K. J., & Gonzalez, J. M. (2020). Biology, strategies, and fresh meat consequences of manipulating the fatty acid composition of meat. *Journal of Animal Science*, 98(2). <https://doi.org/10.1093/jas/skaa033>
- Butler, G., Ali, A. M., Oladokun, S., Wang, J., & Davis, H. (2021). Forage-fed cattle point the way forward for beef? *Future Foods*, 3, 100012. <https://doi.org/10.1016/j.fufo.2021.100012>
- Cabrera, M., & Saadoun, A. (2014). An overview of the nutritional value of beef and lamb meat from South America. *Meat Science*, 98(3), 435–444. <https://doi.org/10.1016/j.meatsci.2014.06.033>

- Cama-Moncunill, R., Moloney, A., Röhrle, F., Luciano, G., & Monahan, F. (2021). Canonical discriminant analysis of the fatty acid profile of muscle to authenticate beef from grass-fed and other beef production systems: Model development and validation. *Food Control*, *122*, 107820. <https://doi.org/10.1016/j.foodcont.2020.107820>
- Carrillo, J. A., He, Y., Li, Y., Liu, J., Erdman, R. A., Sonstegard, T. S., & Song, J. (2016). Integrated metabolomic and transcriptome analyses reveal finishing forage affects metabolic pathways related to beef quality and animal welfare. *Scientific Reports*, *6*(1). <https://doi.org/10.1038/srep25948>
- Casperson, S. L., Conrad, Z., Raatz, S. K., Derner, J., Roemmich, J. N., Jahns, L., & Picklo, M. J. (2020). Impact of beef consumption on saturated fat intake in the United States adult population: Insights from modeling the influences of bovine genetics and nutrition. *Meat Science*, *169*, 108225. <https://doi.org/10.1016/j.meatsci.2020.108225>
- Chail, A., Legako, J. F., Pitcher, L. R., Griggs, T. C., Ward, R. E., Martini, S., & MacAdam, J. W. (2016). Legume finishing provides beef with positive human dietary fatty acid ratios and consumer preference comparable with grain-finished beef¹. *Journal of Animal Science*, *94*(5), 2184–2197. <https://doi.org/10.2527/jas.2015-0241>
- Chail, A., Legako, J. F., Pitcher, L. R., Ward, R. E., Martini, S., & MacAdam, J. W. (2017). Consumer sensory evaluation and chemical composition of beef gluteus medius and triceps brachii steaks from cattle finished on forage or concentrate diets¹. *Journal of Animal Science*, *95*(4), 1553–1564. <https://doi.org/10.2527/jas.2016.1150>
- Cherfaoui, M., Durand, D., Bonnet, M., Bernard, L., Bauchart, D., Ortigues-Marty, I., & Gruffat, D. (2013). A grass-based diet favours muscle n-3 long-chain PUFA deposition without modifying gene expression of proteins involved in their synthesis or uptake in Charolais steers. *Animal*, *7*(11), 1833–1840. <https://doi.org/10.1017/s1751731113001432>
- da Costa, A. S., Bessa, R. J., Pires, V. M., Rolo, E., Pinto, R. M., Andrade Fontes, C., & Prates, J. A. (2014). Is hepatic lipid metabolism of beef cattle influenced by breed and dietary silage level? *BMC Veterinary Research*, *10*(1), 65. <https://doi.org/10.1186/1746-6148-10-65>
- da Costa, A. S., Pires, V. M., Fontes, C. M., & Mestre Prates, J. A. (2013). Expression of genes controlling fat deposition in two genetically diverse beef cattle breeds fed high or low silage diets. *BMC Veterinary Research*, *9*(1), 118. <https://doi.org/10.1186/1746-6148-9-118>
- Frank, D., Ball, A., Hughes, J., Krishnamurthy, R., Piyasiri, U., Stark, J., Watkins, P., & Warner, R. (2016). Sensory and Flavor Chemistry Characteristics of Australian Beef: Influence of Intramuscular Fat, Feed, and Breed. *Journal of Agricultural and Food Chemistry*, *64*(21), 4299–4311. <https://doi.org/10.1021/acs.jafc.6b00160>
- Freitas, A. D., Lobato, J., Cardoso, L., Tarouco, J., Vieira, R., Dillenburg, D., & Castro, I. (2014). Nutritional composition of the meat of Hereford and Braford steers finished on pastures or in a feedlot in southern Brazil. *Meat Science*, *96*(1), 353–360. <https://doi.org/10.1016/j.meatsci.2013.07.021>

- Fruet, A., Trombetta, F., Stefanello, F., Speroni, C., Donadel, J., de Souza, A., Rosado Júnior, A., Tonetto, C., Wagner, R., de Mello, A., & Nörnberg, J. (2018). Effects of feeding legume-grass pasture and different concentrate levels on fatty acid profile, volatile compounds, and off-flavor of the *M. longissimus thoracis*. *Meat Science*, *140*, 112–118. <https://doi.org/10.1016/j.meatsci.2018.03.008>
- Greenwood, P. L., Siddell, J. P., Walmsley, B. J., Geesink, G. H., Pethick, D. W., & McPhee, M. J. (2015). Postweaning substitution of grazed forage with a high-energy concentrate has variable long-term effects on subcutaneous fat and marbling in *Bos taurus* genotypes 1. *Journal of Animal Science*, *93*(8), 4132–4143. <https://doi.org/10.2527/jas.2015-8962>
- Hall, N., Schönfeldt, H., & Pretorius, B. (2016). Fatty acids in beef from grain- and grass-fed cattle: the unique South African scenario. *South African Journal of Clinical Nutrition*, *29*(2), 55–62. <https://doi.org/10.1080/16070658.2016.1216359>
- Haskins, C. P., Henderson, G., & Champ, C. E. (2018). Meat, eggs, full-fat dairy, and nutritional boogeymen: Does the way in which animals are raised affect health differently in humans? *Critical Reviews in Food Science and Nutrition*, *59*(17), 2709–2719. <https://doi.org/10.1080/10408398.2018.1465888>
- He, M. L., Stanford, K., Dugan, M. E. R., Marquess, L., & McAllister, T. A. (2020). Association of leptin genotype with growth performance, adipocyte cellularity, meat quality, and fatty acid profile in beef steers fed flaxseed or high-oleate sunflower seed diets with or without triticale dried distiller's grains. *Journal of Animal Science*, *98*(4). <https://doi.org/10.1093/jas/skaa104>
- Horcada, A., López, A., Polvillo, O., Pino, R., Cubiles-de-la-Vega, D., Tejerina, D., & García-Torres, S. (2018). Fatty acid profile as a tool to trace the origin of beef in pasture- and grain-fed young bulls of Retinta breed. *Spanish Journal of Agricultural Research*, *15*(4), e0607. <https://doi.org/10.5424/sjar/2017154-11032>
- Horcada, A., Polvillo, O., Juárez, M., Avilés, C., Martínez, A., & Peña, F. (2016). Influence of feeding system (concentrate and total mixed ration) on fatty acid profiles of beef from three lean cattle breeds. *Journal of Food Composition and Analysis*, *49*, 110–116. <https://doi.org/10.1016/j.jfca.2016.04.008>
- Humada, M., Sañudo, C., & Serrano, E. (2014). Chemical composition, vitamin E content, lipid oxidation, colour and cooking losses in meat from Tudanca bulls finished on semi-extensive or intensive systems and slaughtered at 12 or 14 months. *Meat Science*, *96*(2), 908–915. <https://doi.org/10.1016/j.meatsci.2013.10.004>
- Hwang, Y. H., & Joo, S. T. (2017). Fatty Acid Profiles, Meat Quality, and Sensory Palatability of Grain-fed and Grass-fed Beef from Hanwoo, American, and Australian Crossbred Cattle. *Korean Journal for Food Science of Animal Resources*, *37*(2), 153–161. <https://doi.org/10.5851/kosfa.2017.37.2.153>

- Kamihiro, S., Stergiadis, S., Leifert, C., Eyre, M., & Butler, G. (2015). Meat quality and health implications of organic and conventional beef production. *Meat Science*, *100*, 306–318. <https://doi.org/10.1016/j.meatsci.2014.10.015>
- Lafreniere, C., Berthiaume, R., Giesen, L., Campbell, C., Pivotto-Baird, L., & Mandell, I. (2021). Effects of forage finishing methods with alfalfa on cattle growth performance and beef carcass characteristics, eating quality, and nutrient composition. *Canadian Journal of Animal Science*, *101*(1), 30–48. <https://doi.org/10.1139/cjas-2019-0121>
- ŁOzicki, A., & Dymnicka, M. (2014). Effect of beef from grass or maize silage- and concentrate-fed cattle on lipid metabolism and antioxidative status of rats. *Journal of Animal and Feed Sciences*, *23*(1), 45–51. <https://doi.org/10.22358/jafs/65715/2014>
- MacKintosh, S. B., Richardson, I., Kim, E. J., Dannenberger, D., Coulmier, D., & Scollan, N. D. (2017). Addition of an extract of lucerne (*Medicago sativa* L.) to cattle diets – Effects on fatty acid profile, meat quality and eating quality of the M. longissimus muscle. *Meat Science*, *130*, 69–80. <https://doi.org/10.1016/j.meatsci.2017.03.011>
- Mapiye, C., Aalhus, J., Turner, T., Vahmani, P., Baron, V., McAllister, T., Block, H., Uttaro, B., & Dugan, M. (2014). Inclusion of sunflower seed and wheat dried distillers' grains with solubles in a red clover silage-based diet enhances steers performance, meat quality and fatty acid profiles. *Animal*, *8*(12), 1999–2010. <https://doi.org/10.1017/s1751731114001955>
- Mezgebo, G. B., Monahan, F. J., McGee, M., O'Riordan, E. G., Richardson, I. R., Brunton, N. P., & Moloney, A. P. (2017). Fatty acid, volatile and sensory characteristics of beef as affected by grass silage or pasture in the bovine diet. *Food Chemistry*, *235*, 86–97. <https://doi.org/10.1016/j.foodchem.2017.05.025>
- Mezgebo, G., Monahan, F., McGee, M., O'Riordan, E., Picard, B., Richardson, R., & Moloney, A. (2017). Biochemical and organoleptic characteristics of muscle from early and late maturing bulls in different production systems. *Animal*, *11*(9), 1636–1644. <https://doi.org/10.1017/s175173111600272x>
- Moloney, A., Mooney, M., Kerry, J., Stanton, C., & O'Kiely, P. (2013). Colour of fat, and colour, fatty acid composition and sensory characteristics of muscle from heifers offered alternative forages to grass silage in a finishing ration. *Meat Science*, *95*(3), 608–615. <https://doi.org/10.1016/j.meatsci.2013.05.030>
- Moran, L., O'Sullivan, M., Kerry, J., Picard, B., McGee, M., O'Riordan, E., & Moloney, A. (2017). Effect of a grazing period prior to finishing on a high concentrate diet on meat quality from bulls and steers. *Meat Science*, *125*, 76–83. <https://doi.org/10.1016/j.meatsci.2016.11.021>
- Patel, M., Sonesson, U., & Hesse, A. (2017). Upgrading plant amino acids through cattle to improve the nutritional value for humans: effects of different production systems. *Animal*, *11*(3), 519–528. <https://doi.org/10.1017/s1751731116001610>

- Patino, H., Medeiros, F., Pereira, C., Swanson, K., & McManus, C. (2015). Productive performance, meat quality and fatty acid profile of steers finished in confinement or supplemented at pasture. *Animal*, 9(6), 966–972. <https://doi.org/10.1017/s1751731115000105>
- Ribas-Agustí, A., Díaz, I., Sárraga, C., García-Regueiro, J. A., & Castellari, M. (2019). Nutritional properties of organic and conventional beef meat at retail. *Journal of the Science of Food and Agriculture*, 99(9), 4218–4225. <https://doi.org/10.1002/jsfa.9652>
- Ripoll, G., Blanco, M., Albertí, P., Panea, B., Joy, M., & Casasús, I. (2013). Effect of two Spanish breeds and diet on beef quality including consumer preferences. *Journal of the Science of Food and Agriculture*, 94(5), 983–992. <https://doi.org/10.1002/jsfa.6348>
- Rosa, H. J. D., Rego, O. A., Silva, C. C. G., Alves, S. P., Alfaia, C. M. M., Prates, J. A. M., & Bessa, R. J. B. (2014). Effect of corn supplementation of grass finishing of Holstein bulls on fatty acid composition of meat lipids1. *Journal of Animal Science*, 92(8), 3701–3714. <https://doi.org/10.2527/jas.2013-6982>
- Santos-Silva, J., Alves, S. P., Francisco, A., Portugal, A. P., Almeida, J., Fialho, L., Jerónimo, E., & Bessa, R. J. (2020). Effects of a high-fibre and low-starch diet in growth performance, carcass and meat quality of young Alentejana breed bulls. *Meat Science*, 168, 108191. <https://doi.org/10.1016/j.meatsci.2020.108191>
- Scollan, N. D., Dannenberger, D., Nuernberg, K., Richardson, I., MacKintosh, S., Hocquette, J. F., & Moloney, A. P. (2014). Enhancing the nutritional and health value of beef lipids and their relationship with meat quality. *Meat Science*, 97(3), 384–394. <https://doi.org/10.1016/j.meatsci.2014.02.015>
- Shibata, M., Hikino, Y., Imanari, M., & Matsumoto, K. (2019). Comprehensive evaluation of growth performance and meat characteristics of a fattening system combining grazing with feeding rice whole-crop silage in Japanese Black steers. *Animal Science Journal*, 90(4), 504–512. <https://doi.org/10.1111/asj.13176>
- Shibata, M., Hikino, Y., & Matsumoto, K. (2018). Influence of feeding a grass hay diet during the early stage of the fattening period on growth performance, carcass characteristics, and meat production in Japanese Black steers. *Animal Science Journal*, 90(2), 196–204. <https://doi.org/10.1111/asj.13139>
- Siphambili, S., Moloney, A., O’Riordan, E., McGee, M., & Monahan, F. (2020). The effects of graded levels of concentrate supplementation on colour and lipid stability of beef from pasture finished late-maturing bulls. *Animal*, 14(3), 656–666. <https://doi.org/10.1017/s1751731119002313>
- Smith, S. B., Lunt, D. K., Smith, D. R., & Walzem, R. L. (2020). Producing high-oleic acid beef and the impact of ground beef consumption on risk factors for cardiovascular disease: A review. *Meat Science*, 163, 108076. <https://doi.org/10.1016/j.meatsci.2020.108076>

- Soliman, L. C., Andrucson, E. M., Donkor, K. K., Church, J. S., & Cinel, B. (2015). Determination of Fatty Acids in Beef by Liquid Chromatography-Electrospray Ionization Tandem Mass Spectrometry. *Food Analytical Methods*, 9(3), 630–637. <https://doi.org/10.1007/s12161-015-0229-0>
- Soulat, J., Picard, B., Léger, S., & Monteils, V. (2016). Prediction of beef carcass and meat traits from rearing factors in young bulls and cull cows. *Journal of Animal Science*, 94(4), 1712–1726. <https://doi.org/10.2527/jas.2015-0164>
- Średnicka-Tober, D., Barański, M., Seal, C., Sanderson, R., Benbrook, C., Steinshamn, H., Gromadzka-Ostrowska, J., Rembiałkowska, E., Skwarło-Sońta, K., Eyre, M., Cozzi, G., Krogh Larsen, M., Jordon, T., Niggli, U., Sakowski, T., Calder, P. C., Burdge, G. C., Sotiraki, S., Stefanakis, A., . . . Leifert, C. (2016). Composition differences between organic and conventional meat: a systematic literature review and meta-analysis. *British Journal of Nutrition*, 115(6), 994–1011. <https://doi.org/10.1017/s0007114515005073>
- Tansawat, R., Maughan, C. A. J., Ward, R. E., Martini, S., & Cornforth, D. P. (2012). Chemical characterisation of pasture- and grain-fed beef related to meat quality and flavour attributes. *International Journal of Food Science & Technology*, 48(3), 484–495. <https://doi.org/10.1111/j.1365-2621.2012.03209.x>
- van Elswyk, M. E., & McNeill, S. H. (2014). Impact of grass/forage feeding versus grain finishing on beef nutrients and sensory quality: The U.S. experience. *Meat Science*, 96(1), 535–540. <https://doi.org/10.1016/j.meatsci.2013.08.010>
- van Vliet, S., Provenza, F. D., & Kronberg, S. L. (2021). Health-Promoting Phytonutrients Are Higher in Grass-Fed Meat and Milk. *Frontiers in Sustainable Food Systems*, 4. <https://doi.org/10.3389/fsufs.2020.555426>
- Venkata Reddy, B., Sivakumar, A. S., Jeong, D. W., Woo, Y. B., Park, S. J., Lee, S. Y., Byun, J. Y., Kim, C. H., Cho, S. H., & Hwang, I. (2014). Beef quality traits of heifer in comparison with steer, bull and cow at various feeding environments. *Animal Science Journal*, 86(1), 1–16. <https://doi.org/10.1111/asj.12266>
- Wright, A. M., Andrae, J. G., Rosso, C. F., Miller, M. C., Pavan, E., Bridges, W., & Duckett, S. K. (2015). Effect of forage type with or without corn supplementation on animal performance, beef fatty acid composition, and palatability¹². *Journal of Animal Science*, 93(10), 5047–5058. <https://doi.org/10.2527/jas.2015-8939>

Bison

- Daley, C. A., Abbott, A., Doyle, P. S., Nader, G. A., & Larson, S. (2010). A review of fatty acid profiles and antioxidant content in grass-fed and grain-fed beef. *Nutrition Journal*, 9(1). <https://doi.org/10.1186/1475-2891-9-10>
- DRISKELL, J. A., MARCHELLO, M. J., GIRAUD, D. W., & SULAEMAN, A. (2004). VITAMIN AND SELENIUM CONTENT OF RIBEYE CUTS FROM GRASS- AND

GRAIN-FINISHED BISON OF THE SAME HERD. *Journal of Food Quality*, 27(5), 388–398. <https://doi.org/10.1111/j.1745-4557.2004.00680.x>

Jutzeler Van Wijlen, R. P., & Colombani, P. C. (2010). Grass-based ruminant production methods and human bioconversion of vaccenic acid with estimations of maximal dietary intake of conjugated linoleic acids. *International Dairy Journal*, 20(7), 433–448. <https://doi.org/10.1016/j.idairyj.2010.01.008>

Lulka, D. (2006). Grass or Grain? Assessing the Nature of the US Bison Industry. *Sociologia Ruralis*, 46(3), 173–191. <https://doi.org/10.1111/j.1467-9523.2006.00413.x>

McDaniel, J., Askew, W., Bennett, D., Mihalopoulos, J., Anantharaman, S., Fjeldstad, A. S., Rule, D. C., Nanjee, N. M., Harris, R. A., & Richardson, R. S. (2013). Bison meat has a lower atherogenic risk than beef in healthy men. *Nutrition Research*, 33(4), 293–302. <https://doi.org/10.1016/j.nutres.2013.01.007>

Mir, S., Ahmad, H., Ganai, A., Lone, S., Ganaie, B., & Malik, T. (2018). Role of Pastures in Value Addition of Animal Products: A Review. *International Journal of Livestock Research*, 8(5), 35. <https://doi.org/10.5455/ijlr.20170818115235>

Rule, D. C., Broughton, K. S., Shellito, S. M., & Maiorano, G. (2002). Comparison of muscle fatty acid profiles and cholesterol concentrations of bison, beef cattle, elk, and chicken1. *Journal of Animal Science*, 80(5), 1202–1211. <https://doi.org/10.2527/2002.8051202x>

Turner, T. D., Pilfold, J. L., Jensen, J., Prema, D., Donkor, K. K., van Hamme, J. D., Cinel, B., Galbraith, J. K., & Church, J. S. (2014). Fatty Acid Profiles of Western Canadian Bison (Bison bison) Meat. *Journal of Food Research*, 3(6), 146. <https://doi.org/10.5539/jfr.v3n6p146>

Vahmani, P., Ponnampalam, E. N., Kraft, J., Mapiye, C., Bermingham, E. N., Watkins, P. J., Proctor, S. D., & Dugan, M. E. (2020). Bioactivity and health effects of ruminant meat lipids. Invited Review. *Meat Science*, 165, 108114. <https://doi.org/10.1016/j.meatsci.2020.108114>

van Vliet, S., Provenza, F. D., & Kronberg, S. L. (2021). Health-Promoting Phytonutrients Are Higher in Grass-Fed Meat and Milk. *Frontiers in Sustainable Food Systems*, 4. <https://doi.org/10.3389/fsufs.2020.555426>

Dairy cows

Akbaridoust, G., Plozza, T., Trenerry, V. C., Wales, W. J., Auldust, M. J., & Ajlouni, S. (2015). Influence of pasture-based feeding systems on fatty acids, organic acids and volatile organic flavour compounds in yoghurt. *Journal of Dairy Research*, 82(3), 279–286. <https://doi.org/10.1017/s0022029915000357>

Barca, J., Carriquiry, M., Olazabal, L., Fajardo, M., Chilibroste, P., & Meikle, A. (2017). Milk fatty acid profile from cows fed with mixed rations and different access time to pastureland

- during early lactation. *Journal of Animal Physiology and Animal Nutrition*, 102(3), 620–629. <https://doi.org/10.1111/jpn.12826>
- Bittante, G., Cecchinato, A., & Schiavon, S. (2018). Dairy system, parity, and lactation stage affect enteric methane production, yield, and intensity per kilogram of milk and cheese predicted from gas chromatography fatty acids. *Journal of Dairy Science*, 101(2), 1752–1766. <https://doi.org/10.3168/jds.2017-13472>
- Butler, G., Stergiadis, S., Chatzidimitriou, E., Franceschin, E., Davis, H. R., Leifert, C., & Steinshamn, H. (2019). Differing responses in milk composition from introducing rapeseed and naked oats to conventional and organic dairy diets. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-44567-8>
- Cavaliere, G., Trinchese, G., Musco, N., Infascelli, F., de Filippo, C., Mastellone, V., Morittu, V. M., Lombardi, P., Tudisco, R., Grossi, M., Monda, V., Cutrignelli, M. I., Messina, A., Calabrò, S., Moni, H. B., Stradella, L., Messina, G., Monda, M., Crispino, M., & Mollica, M. P. (2018). Milk from cows fed a diet with a high forage:concentrate ratio improves inflammatory state, oxidative stress, and mitochondrial function in rats. *Journal of Dairy Science*, 101(3), 1843–1851. <https://doi.org/10.3168/jds.2017-13550>
- Coppa, M., Chassaing, C., Sibra, C., Cornu, A., Verbič, J., Golecký, J., Engel, E., Ratel, J., Boudon, A., Ferlay, A., & Martin, B. (2019). Forage system is the key driver of mountain milk specificity. *Journal of Dairy Science*, 102(11), 10483–10499. <https://doi.org/10.3168/jds.2019-16726>
- Davis, H., Chatzidimitriou, E., Leifert, C., & Butler, G. (2020). Evidence That Forage-Fed Cows Can Enhance Milk Quality. *Sustainability*, 12(9), 3688. <https://doi.org/10.3390/su12093688>
- Faulkner, H., O’Callaghan, T. F., McAuliffe, S., Hennessy, D., Stanton, C., O’Sullivan, M. G., Kerry, J. P., & Kilcawley, K. N. (2018). Effect of different forage types on the volatile and sensory properties of bovine milk. *Journal of Dairy Science*, 101(2), 1034–1047. <https://doi.org/10.3168/jds.2017-13141>
- Frétin, M., Martin, B., Buchin, S., Desserre, B., Lavigne, R., Tixier, E., Cirié, C., Bord, C., Montel, M. C., Delbès, C., & Ferlay, A. (2019). Milk fat composition modifies the texture and appearance of Cantal-type cheeses but not their flavor. *Journal of Dairy Science*, 102(2), 1131–1143. <https://doi.org/10.3168/jds.2018-15534>
- Gómez-Mascaraque, L., Kilcawley, K., Hennessy, D., Tobin, J., & O’Callaghan, T. (2020). Raman spectroscopy: A rapid method to assess the effects of pasture feeding on the nutritional quality of butter. *Journal of Dairy Science*, 103(10), 8721–8731. <https://doi.org/10.3168/jds.2020-18716>
- Haskins, C. P., Henderson, G., & Champ, C. E. (2018). Meat, eggs, full-fat dairy, and nutritional boogeymen: Does the way in which animals are raised affect health differently in humans? *Critical Reviews in Food Science and Nutrition*, 59(17), 2709–2719. <https://doi.org/10.1080/10408398.2018.1465888>

- Hegazy, M., Elsayed, N. M., Ali, H. M., Hassan, H. G., & Rashed, L. (2019). Diabetes Mellitus, Nonalcoholic Fatty Liver Disease, and Conjugated Linoleic Acid (Omega 6): What Is the Link? *Journal of Diabetes Research*, 2019, 1–7. <https://doi.org/10.1155/2019/5267025>
- Hur, S. J., Kim, H. S., Bahk, Y. Y., & Park, Y. (2017). Overview of conjugated linoleic acid formation and accumulation in animal products. *Livestock Science*, 195, 105–111. <https://doi.org/10.1016/j.livsci.2016.11.016>
- Jaakamo, M. J., Luukkonen, T. J., Kairenius, P. K., Bayat, A. R., Ahvenjärvi, S. A., Tupasela, T. M., Vilkki, J. H., Shingfield, K. J., & Leskinen, H. M. (2019). The effect of dietary forage to concentrate ratio and forage type on milk fatty acid composition and milk fat globule size of lactating cows. *Journal of Dairy Science*, 102(10), 8825–8838. <https://doi.org/10.3168/jds.2018-15833>
- Khiaosa-ard, R., Kreuzer, M., & Leiber, F. (2015). Apparent recovery of C18 polyunsaturated fatty acids from feed in cow milk: A meta-analysis of the importance of dietary fatty acids and feeding regimens in diets without fat supplementation. *Journal of Dairy Science*, 98(9), 6399–6414. <https://doi.org/10.3168/jds.2015-9459>
- Livingstone, K., Humphries, D., Kirton, P., Kliem, K., Givens, D., & Reynolds, C. (2015). Effects of forage type and extruded linseed supplementation on methane production and milk fatty acid composition of lactating dairy cows. *Journal of Dairy Science*, 98(6), 4000–4011. <https://doi.org/10.3168/jds.2014-8987>
- Magan, J. B., O’Callaghan, T. F., Zheng, J., Zhang, L., Mandal, R., Hennessy, D., Fenelon, M. A., Wishart, D. S., Kelly, A. L., & McCarthy, N. A. (2020). Effect of Diet on the Vitamin B Profile of Bovine Milk-Based Protein Ingredients. *Foods*, 9(5), 578. <https://doi.org/10.3390/foods9050578>
- Mendoza, A., Cajarville, C., & Repetto, J. (2016). Short communication: Intake, milk production, and milk fatty acid profile of dairy cows fed diets combining fresh forage with a total mixed ration. *Journal of Dairy Science*, 99(3), 1938–1944. <https://doi.org/10.3168/jds.2015-10257>
- Mesilati-Stahy, R., Moallem, U., Magen, Y., & Argov-Argaman, N. (2015). Altered concentrate to forage ratio in cows ration enhanced bioproduction of specific size subpopulation of milk fat globules. *Food Chemistry*, 179, 199–205. <https://doi.org/10.1016/j.foodchem.2015.01.138>
- Mir, S., Ahmad, H., Ganai, A., Lone, S., Ganaie, B., & Malik, T. (2018). Role of Pastures in Value Addition of Animal Products: A Review. *International Journal of Livestock Research*, 8(5), 35. <https://doi.org/10.5455/ijlr.20170818115235>
- Mitani, T., Kobayashi, K., Ueda, K., & Kondo, S. (2015). Discrimination of “grazing milk” using milk fatty acid profile in the grassland dairy area in Hokkaido. *Animal Science Journal*, 87(2), 233–241. <https://doi.org/10.1111/asj.12422>

- O'Callaghan, T. F., Faulkner, H., McAuliffe, S., O'Sullivan, M. G., Hennessy, D., Dillon, P., Kilcawley, K. N., Stanton, C., & Ross, R. (2016). Quality characteristics, chemical composition, and sensory properties of butter from cows on pasture versus indoor feeding systems. *Journal of Dairy Science*, 99(12), 9441–9460. <https://doi.org/10.3168/jds.2016-11271>
- O'Callaghan, T. F., Hennessy, D., McAuliffe, S., Kilcawley, K. N., O'Donovan, M., Dillon, P., Ross, R., & Stanton, C. (2016). Effect of pasture versus indoor feeding systems on raw milk composition and quality over an entire lactation. *Journal of Dairy Science*, 99(12), 9424–9440. <https://doi.org/10.3168/jds.2016-10985>
- Pastorini, M., Pomiés, N., Repetto, J., Mendoza, A., & Cajarville, C. (2019). Productive performance and digestive response of dairy cows fed different diets combining a total mixed ration and fresh forage. *Journal of Dairy Science*, 102(5), 4118–4130. <https://doi.org/10.3168/jds.2018-15389>
- Payling, L. M., Juniper, D. T., Drake, C., Rymer, C., & Givens, D. I. (2015). Effect of milk type and processing on iodine concentration of organic and conventional winter milk at retail: Implications for nutrition. *Food Chemistry*, 178, 327–330. <https://doi.org/10.1016/j.foodchem.2015.01.091>
- Peira, G., Cortese, D., Lombardi, G., & Bollani, L. (2020). Grass-Fed Milk Perception: Profiling Italian Consumer. *Sustainability*, 12(24), 10348. <https://doi.org/10.3390/su122410348>
- Puppel, K., Sakowski, T., Kuczyńska, B., Grodkowski, G., Gołębiewski, M., Barszczewski, J., Wróbel, B., Budziński, A., Kapusta, A., & Balcerak, M. (2017). Degrees of Antioxidant Protection: A 2-Year Study of the Bioactive Properties of Organic Milk in Poland. *Journal of Food Science*, 82(2), 523–528. <https://doi.org/10.1111/1750-3841.13594>
- Sabikhi, L. (2007). Designer Milk. *Advances in Food and Nutrition Research*, 161–198. [https://doi.org/10.1016/s1043-4526\(07\)53005-6](https://doi.org/10.1016/s1043-4526(07)53005-6)
- Sankarlal, V., Testroet, E., Beitz, D., & Clark, S. (2015). Dried distillers grains with solubles do not always cause late blowing in baby Swiss cheese. *Journal of Dairy Science*, 98(12), 8545–8553. <https://doi.org/10.3168/jds.2015-9983>
- Schulz, F., Westreicher-Kristen, E., Molkentin, J., Knappstein, K., & Susenbeth, A. (2018). Effect of replacing maize silage with red clover silage in the diet on milk fatty acid composition in cows. *Journal of Dairy Science*, 101(8), 7156–7167. <https://doi.org/10.3168/jds.2017-14335>
- Schwendel, B. H., Wester, T. J., Morel, P. C., Fong, B., Tavendale, M. H., Deadman, C., Shadbolt, N. M., & Otter, D. E. (2017). Pasture feeding conventional cows removes differences between organic and conventionally produced milk. *Food Chemistry*, 229, 805–813. <https://doi.org/10.1016/j.foodchem.2017.02.104>

- Segato, S., Galaverna, G., Contiero, B., Berzaghi, P., Caligiani, A., Marseglia, A., & Cozzi, G. (2017). Identification of Lipid Biomarkers To Discriminate between the Different Production Systems for Asiago PDO Cheese. *Journal of Agricultural and Food Chemistry*, 65(45), 9887–9892. <https://doi.org/10.1021/acs.jafc.7b03629>
- Średnicka-Tober, D., Barański, M., Seal, C. J., Sanderson, R., Benbrook, C., Steinshamn, H., Gromadzka-Ostrowska, J., Rembiałkowska, E., Skwarło-Sońta, K., Eyre, M., Cozzi, G., Larsen, M. K., Jordon, T., Niggli, U., Sakowski, T., Calder, P. C., Burdge, G. C., Sotiraki, S., Stefanakis, A., . . . Leifert, C. (2016). Higher PUFA and n-3 PUFA, conjugated linoleic acid, α -tocopherol and iron, but lower iodine and selenium concentrations in organic milk: a systematic literature review and meta- and redundancy analyses. *British Journal of Nutrition*, 115(6), 1043–1060. <https://doi.org/10.1017/s0007114516000349>
- Stanton, C., Mills, S., Ryan, A., di Gioia, D., & Ross, R. (2021). Influence of pasture feeding on milk and meat products in terms of human health and product quality. *Irish Journal of Agricultural and Food Research*. Published. <https://doi.org/10.15212/ijafr-2020-0104>
- Thomson, A., Humphries, D., Kliem, K., Dittmann, M., & Reynolds, C. (2017). Effects of replacing maize silage with lucerne silage and lucerne silage chop length on rumen function and milk fatty acid composition. *Journal of Dairy Science*, 100(9), 7127–7138. <https://doi.org/10.3168/jds.2017-12914>
- Umaraw, P. (2015). Designer Milk - A Milk of Intrinsic Health Benefit: A Review. *Journal of Food Processing & Technology*, 06(03). <https://doi.org/10.4172/2157-7110.1000426>
- Unger, A. L., Bourne, D. E., Walsh, H., & Kraft, J. (2020). Fatty Acid Content of Retail Cow's Milk in the Northeastern United States—What's in It for the Consumer? *Journal of Agricultural and Food Chemistry*, 68(14), 4268–4276. <https://doi.org/10.1021/acs.jafc.9b07390>
- van Vliet, S., Provenza, F. D., & Kronberg, S. L. (2021). Health-Promoting Phytonutrients Are Higher in Grass-Fed Meat and Milk. *Frontiers in Sustainable Food Systems*, 4. <https://doi.org/10.3389/fsufs.2020.555426>
- Vanbergue, E., Delaby, L., Peyraud, J., Colette, S., Gallard, Y., & Hurtaud, C. (2017). Effects of breed, feeding system, and lactation stage on milk fat characteristics and spontaneous lipolysis in dairy cows. *Journal of Dairy Science*, 100(6), 4623–4636. <https://doi.org/10.3168/jds.2016-12094>
- Wilkinson, J., & Lee, M. (2018). Review: Use of human-edible animal feeds by ruminant livestock. *Animal*, 12(8), 1735–1743. <https://doi.org/10.1017/s175173111700218x>
- Yang, Y., Ferreira, G., Teets, C., Corl, B., Thomason, W., & Griffey, C. (2018). Effects of feeding hulled and hull-less barley with low- and high-forage diets on lactation performance, nutrient digestibility, and milk fatty acid composition of lactating dairy cows. *Journal of Dairy Science*, 101(4), 3036–3043. <https://doi.org/10.3168/jds.2017-14082>

Zendri, F., Ramanzin, M., Cipolat-Gotet, C., & Sturaro, E. (2016). Variation of milk coagulation properties, cheese yield, and nutrients recovery in curd of cows of different breeds before, during and after transhumance to highland summer pastures. *Journal of Dairy Research*, 84(1), 39–48. <https://doi.org/10.1017/s0022029916000583>

Laying Hens

Anderson, K. (2011). Comparison of fatty acid, cholesterol, and vitamin A and E composition in eggs from hens housed in conventional cage and range production facilities. *Poultry Science*, 90(7), 1600–1608. <https://doi.org/10.3382/ps.2010-01289>

Bryden, W. L., Li, X., Ruhnke, I., Zhang, D., & Shini, S. (2021). Nutrition, feeding and laying hen welfare. *Animal Production Science*, 61(10), 893. <https://doi.org/10.1071/an20396>

Dunlop, E., Cunningham, J., Sherriff, J., Lucas, R., Greenfield, H., Arcot, J., Strobel, N., & Black, L. (2017). Vitamin D3 and 25-Hydroxyvitamin D3 Content of Retail White Fish and Eggs in Australia. *Nutrients*, 9(7), 647. <https://doi.org/10.3390/nu9070647>

Grela, E., Knaga, S., Winiarska-Mieczan, A., & Zięba, G. (2020). Effects of dietary alfalfa protein concentrate supplementation on performance, egg quality, and fatty acid composition of raw, freeze-dried, and hard-boiled eggs from Polbar laying hens. *Poultry Science*, 99(4), 2256–2265. <https://doi.org/10.1016/j.psj.2019.11.030>

Guo, J., Kliem, K. E., Lovegrove, J. A., & Givens, D. (2017). Effect of production system, supermarket and purchase date on the vitamin D content of eggs at retail. *Food Chemistry*, 221, 1021–1025. <https://doi.org/10.1016/j.foodchem.2016.11.060>

Haskins, C. P., Henderson, G., & Champ, C. E. (2018). Meat, eggs, full-fat dairy, and nutritional boogeymen: Does the way in which animals are raised affect health differently in humans? *Critical Reviews in Food Science and Nutrition*, 59(17), 2709–2719. <https://doi.org/10.1080/10408398.2018.1465888>

Heflin, L. E., Malheiros, R., Anderson, K. E., Johnson, L. K., & Raatz, S. K. (2018). Mineral content of eggs differs with hen strain, age, and rearing environment. *Poultry Science*, 97(5), 1605–1613. <https://doi.org/10.3382/ps/pey025>

Hisasaga, C., Griffin, S. E., & Tarrant, K. J. (2020). Survey of egg quality in commercially available table eggs. *Poultry Science*, 99(12), 7202–7206. <https://doi.org/10.1016/j.psj.2020.09.049>

Holt, P., Davies, R., Dewulf, J., Gast, R., Huwe, J., Jones, D., Waltman, D., & Willian, K. (2011). The impact of different housing systems on egg safety and quality. *Poultry Science*, 90(1), 251–262. <https://doi.org/10.3382/ps.2010-00794>

Jeni, R. E., Dittoe, D. K., Olson, E. G., Lourenco, J., Seidel, D. S., Ricke, S. C., & Callaway, T. R. (2021). An overview of health challenges in alternative poultry production systems. *Poultry Science*, 100(7), 101173. <https://doi.org/10.1016/j.psj.2021.101173>

- Kühn, J., Schutkowski, A., Kluge, H., Hirche, F., & Stangl, G. I. (2014). Free-range farming: A natural alternative to produce vitamin D-enriched eggs. *Nutrition*, *30*(4), 481–484. <https://doi.org/10.1016/j.nut.2013.10.002>
- Kühn, J., Wassermann, C., Ebschke, S., Schutkowski, A., Thamm, K., Wensch-Dorendorf, M., von Borell, E., & Stangl, G. I. (2019). Feasibility of artificial light regimes to increase the vitamin D content in indoor-laid eggs. *Poultry Science*, *98*(10), 5177–5187. <https://doi.org/10.3382/ps/pez234>
- Lordelo, M., Fernandes, E., Bessa, R., & Alves, S. (2017). Quality of eggs from different laying hen production systems, from indigenous breeds and specialty eggs. *Poultry Science*, *96*(5), 1485–1491. <https://doi.org/10.3382/ps/pew409>
- Mugnai, C., Sossidou, E. N., dal Bosco, A., Ruggeri, S., Mattioli, S., & Castellini, C. (2013). The effects of husbandry system on the grass intake and egg nutritive characteristics of laying hens. *Journal of the Science of Food and Agriculture*, *94*(3), 459–467. <https://doi.org/10.1002/jsfa.6269>
- Nimalaratne, C., & Wu, J. (2015). Hen Egg as an Antioxidant Food Commodity: A Review. *Nutrients*, *7*(10), 8274–8293. <https://doi.org/10.3390/nu7105394>
- Ortiz, D., Lawson, T., Jarrett, R., Ring, A., Scoles, K. L., Hoverman, L., Rocheford, E., Karcher, D. M., & Rocheford, T. (2021). Biofortified orange corn increases xanthophyll density and yolk pigmentation in egg yolks from laying hens. *Poultry Science*, *100*(7), 101117. <https://doi.org/10.1016/j.psj.2021.101117>
- Rakonjac, S., Bogosavljević-Bošković, S., Pavlovski, Z., Škrbić, Z., Dosković, V., Petrović, M., & Petričević, V. (2014a). Laying hen rearing systems: a review of chemical composition and hygienic conditions of eggs. *World's Poultry Science Journal*, *70*(1), 151–164. <https://doi.org/10.1017/s0043933914000130>
- Rakonjac, S., Bogosavljević-Bošković, S., Pavlovski, Z., Škrbić, Z., Dosković, V., Petrović, M., & Petričević, V. (2014b). Laying hen rearing systems: a review of major production results and egg quality traits. *World's Poultry Science Journal*, *70*(1), 93–104. <https://doi.org/10.1017/s0043933914000087>
- Sergin, S., Goeden, T., Krusinski, L., Kesamneni, S., Ali, H., Bitler, C. A., Medina-Meza, I. G., & Fenton, J. I. (2021). Fatty Acid and Antioxidant Composition of Conventional Compared to Pastured Eggs: Characterization of Conjugated Linoleic Acid and Branched Chain Fatty Acid Isomers in Eggs. *ACS Food Science & Technology*, *1*(2), 260–267. <https://doi.org/10.1021/acsfoodscitech.0c00093>
- Sossidou, E., dal Bosco, A., Castellini, C., & Grashorn, M. (2015). Effects of pasture management on poultry welfare and meat quality in organic poultry production systems. *World's Poultry Science Journal*, *71*(2), 375–384. <https://doi.org/10.1017/s0043933915000379>

- Sun, H., Lee, E., Samaraweera, H., Persia, M., & Ahn, D. U. (2013). Effects of increasing concentrations of corn distillers dried grains with solubles on chemical composition and nutrient content of egg. *Poultry Science*, 92(1), 233–242. <https://doi.org/10.3382/ps.2012-02346>
- Trupia, S., Winkler-Moser, J., Guney, A., Beckstead, R., & Chen, C. Y. O. (2016). Nutritional quality of eggs from hens fed distillers dried grains with solubles. *Poultry Science*, 95(11), 2592–2601. <https://doi.org/10.3382/ps/pew142>
- Tufarelli, V., Ragni, M., & Laudadio, V. (2018). Feeding Forage in Poultry: A Promising Alternative for the Future of Production Systems. *Agriculture*, 8(6), 81. <https://doi.org/10.3390/agriculture8060081>

Meat birds

- Almasi, A., Andrassyne, B. G., Milisits, G., Kustosne, P., & Suto, Z. (2015). Effects of different rearing systems on muscle and meat quality traits of slow- and medium-growing male chickens. *British Poultry Science*, 56(3), 320–324. <https://doi.org/10.1080/00071668.2015.1016478>
- Bartlett, J. R. (2015). Comparing the Effects of Conventional and Pastured Poultry Production Systems on Broiler Performance and Meat Quality. *Journal of Agriculture and Life Sciences*, 2(1), 29–36. http://jalsnet.com/journals/Vol_2_No_1_June_2015/5.pdf
- Boz, M. A., Sarica, M., & Yamak, U. S. (2016). Production traits of artificially and naturally hatched geese in intensive and free-range systems – II: slaughter, carcass and meat quality traits. *British Poultry Science*, 58(2), 166–176. <https://doi.org/10.1080/00071668.2016.1261998>
- Chen, X., Jiang, W., Tan, H., Xu, G., Zhang, X., Wei, S., & Wang, X. (2013). Effects of outdoor access on growth performance, carcass composition, and meat characteristics of broiler chickens. *Poultry Science*, 92(2), 435–443. <https://doi.org/10.3382/ps.2012-02360>
- dal Bosco, A., Mugnai, C., Mattioli, S., Rosati, A., Ruggeri, S., Ranucci, D., & Castellini, C. (2016). Transfer of bioactive compounds from pasture to meat in organic free-range chickens. *Poultry Science*, 95(10), 2464–2471. <https://doi.org/10.3382/ps/pev383>
- Dalziel, C. J., Kliem, K. E., & Givens, D. I. (2015). Fat and fatty acid composition of cooked meat from UK retail chickens labelled as from organic and non-organic production systems. *Food Chemistry*, 179, 103–108. <https://doi.org/10.1016/j.foodchem.2015.01.118>
- Geldenhuis, G., Muller, N., & Hoffman, L. C. (2016). The influence of season on the sensory profile of Egyptian goose (*Alopochen aegyptiacus*) meat. *Poultry Science*, 95(9), 2174–2185. <https://doi.org/10.3382/ps/pew135>
- Jaturasitha, S., Chaiwang, N., Kayan, A., & Kreuzer, M. (2016). Nutritional strategies to improve the lipid composition of meat, with emphasis on Thailand and Asia. *Meat Science*, 120, 157–166. <https://doi.org/10.1016/j.meatsci.2016.04.014>

- Jiang, S., Gou, Z., Li, L., Lin, X., & Jiang, Z. (2017). Growth performance, carcass traits and meat quality of yellow-feathered broilers fed graded levels of alfalfa meal with or without wheat. *Animal Science Journal*, 89(3), 561–569. <https://doi.org/10.1111/asj.12968>
- Kim, H. J., Kim, H. J., Jeon, J., Nam, K. C., Shim, K. S., Jung, J. H., Kim, K. S., Choi, Y., Kim, S. H., & Jang, A. (2020). Comparison of the quality characteristics of chicken breast meat from conventional and animal welfare farms under refrigerated storage. *Poultry Science*, 99(3), 1788–1796. <https://doi.org/10.1016/j.psj.2019.12.009>
- Koçer, B., Bozkurt, M., Ege, G., Tüzün, A. E., Konak, R., & Olgun, O. (2018). Effects of a meal feeding regimen and the availability of fresh alfalfa on growth performance and meat and bone quality of broiler genotypes. *British Poultry Science*, 59(3), 318–329. <https://doi.org/10.1080/00071668.2018.1440378>
- Mikulski, D., Celej, J., Jankowski, J., Majewska, T., & Mikulska, M. (2011). Growth Performance, Carcass Traits and Meat Quality of Slower-growing and Fast-growing Chickens Raised with and without Outdoor Access. *Asian-Australasian Journal of Animal Sciences*, 24(10), 1407–1416. <https://doi.org/10.5713/ajas.2011.11038>
- Milićević, D., Vranić, D., Mašić, Z., Parunović, N., Trbović, D., Nedeljković-Trailović, J., & Petrović, Z. (2014). The role of total fats, saturated/unsaturated fatty acids and cholesterol content in chicken meat as cardiovascular risk factors. *Lipids in Health and Disease*, 13(1). <https://doi.org/10.1186/1476-511x-13-42>
- Sales, J. (2014). Effects of access to pasture on performance, carcass composition, and meat quality in broilers: A meta-analysis. *Poultry Science*, 93(6), 1523–1533. <https://doi.org/10.3382/ps.2013-03499>
- Song, Y., Li, Y., Zheng, S., Dai, W., Shen, X., Zhang, Y., Zhao, W., Chang, G., Xu, Q., & Chen, G. (2017). Effects of forage feeding versus grain feeding on the growth performance and meat quality of Yangzhou geese. *British Poultry Science*, 58(4), 397–401. <https://doi.org/10.1080/00071668.2017.1307942>
- Sossidou, E., dal Bosco, A., Castellini, C., & Grashorn, M. (2015). Effects of pasture management on poultry welfare and meat quality in organic poultry production systems. *World's Poultry Science Journal*, 71(2), 375–384. <https://doi.org/10.1017/s0043933915000379>
- Średnicka-Tober, D., Barański, M., Seal, C., Sanderson, R., Benbrook, C., Steinshamn, H., Gromadzka-Ostrowska, J., Rembiałkowska, E., Skwarło-Sońta, K., Eyre, M., Cozzi, G., Krogh Larsen, M., Jordon, T., Niggli, U., Sakowski, T., Calder, P. C., Burdge, G. C., Sotiraki, S., Stefanakis, A., . . . Leifert, C. (2016). Composition differences between organic and conventional meat: a systematic literature review and meta-analysis. *British Journal of Nutrition*, 115(6), 994–1011. <https://doi.org/10.1017/s0007114515005073>
- Stadig, L. M., Rodenburg, T. B., Reubens, B., Aerts, J., Duquenne, B., & Tuytens, F. A. (2016). Effects of free-range access on production parameters and meat quality, composition and

taste in slow-growing broiler chickens. *Poultry Science*, 95(12), 2971–2978.
<https://doi.org/10.3382/ps/pew226>

Tong, H., Cai, J., Lu, J., Wang, Q., Shao, D., & Zou, J. (2015). Effects of outdoor access days on growth performance, carcass yield, meat quality, and lymphoid organ index of a local chicken breed. *Poultry Science*, 94(6), 1115–1121. <https://doi.org/10.3382/ps/pev032>

Tong, H., Wang, Q., Lu, J., Zou, J., Chang, L., & Fu, S. (2014). Effect of free-range days on a local chicken breed: Growth performance, carcass yield, meat quality, and lymphoid organ index. *Poultry Science*, 93(8), 1883–1889. <https://doi.org/10.3382/ps.2013-03470>

Tufarelli, V., Ragni, M., & Laudadio, V. (2018). Feeding Forage in Poultry: A Promising Alternative for the Future of Production Systems. *Agriculture*, 8(6), 81.
<https://doi.org/10.3390/agriculture8060081>

Pigs

Aaslyng, M. D., & Meinert, L. (2017). Meat flavour in pork and beef – From animal to meal. *Meat Science*, 132, 112–117. <https://doi.org/10.1016/j.meatsci.2017.04.012>

Alfaia, C., Lopes, P., Madeira, M., Pestana, J., Coelho, D., Toldrá, F., & Prates, J. (2019). Current feeding strategies to improve pork intramuscular fat content and its nutritional quality. *Advances in Food and Nutrition Research*, 53–94.
<https://doi.org/10.1016/bs.afnr.2019.03.006>

Almeida, J., Bressan, M. C., Santos-Silva, J., Moreira, O., Bettencourt, C., & Gama, L. T. (2018). Physicochemical characteristics and sensory attributes of meat from heavy-weight Iberian and F1 Large White × Landrace pigs finished intensively or in free-range conditions 1. *Journal of Animal Science*, 96(7), 2734–2746. <https://doi.org/10.1093/jas/sky181>

Ayuso, D., González, A., Hernández, F., Peña, F., & Izquierdo, M. (2014). Effect of sex and final fattening on ultrasound and carcass traits in Iberian pigs. *Meat Science*, 96(1), 562–567.
<https://doi.org/10.1016/j.meatsci.2013.08.018>

AYUSO, D., GONZÁLEZ, A., PEÑA, F., & IZQUIERDO, M. (2018). Changes in adipose cells of Longissimus dorsi muscle in Iberian pigs raised under extensive conditions. *Anais Da Academia Brasileira de Ciências*, 90(1), 247–253. <https://doi.org/10.1590/0001-3765201820150567>

Baer, A. A., & Dilger, A. C. (2014). Effect of fat quality on sausage processing, texture, and sensory characteristics. *Meat Science*, 96(3), 1242–1249.
<https://doi.org/10.1016/j.meatsci.2013.11.001>

Bergstrom, J. R., Nelssen, J. L., Tokach, M. D., Dritz, S. S., Goodband, R. D., & DeRouchey, J. M. (2014). The effects of feeder design and dietary dried distillers' grains with solubles on the performance and carcass characteristics of finishing pigs 1,2. *Journal of Animal Science*, 92(8), 3591–3597. <https://doi.org/10.2527/jas.2014-7686>

- Burnett, D. D., Legako, J. F., Phelps, K. J., & Gonzalez, J. M. (2020). Biology, strategies, and fresh meat consequences of manipulating the fatty acid composition of meat. *Journal of Animal Science*, 98(2). <https://doi.org/10.1093/jas/skaa033>
- Fang, J., Cao, Y., Matsuzaki, M., Suzuki, H., & Kimura, H. (2016). Effects of apple pomace-mixed silage on growth performance and meat quality in finishing pigs. *Animal Science Journal*, 87(12), 1516–1521. <https://doi.org/10.1111/asj.12601>
- Harris, E., Mellencamp, M., Johnston, L., Cox, R., & Shurson, G. (2017). Effect of time interval between the second Improvest® dose and slaughter and corn dried distillers grains with solubles feeding strategies on carcass composition, primal cutout, and pork quality of immunologically castrated pigs. *Meat Science*, 127, 13–20. <https://doi.org/10.1016/j.meatsci.2017.01.003>
- Harris, E., Mellencamp, M., Johnston, L., Cox, R., & Shurson, G. (2018). Effectiveness of different corn dried distillers grains with solubles feeding strategies and increasing the time intervals between the second Improvest dose and slaughter of immunologically castrated pigs on belly and pork fat quality. *Meat Science*, 135, 62–73. <https://doi.org/10.1016/j.meatsci.2017.08.025>
- Holinger, M., Früh, B., Stoll, P., Pedan, V., Kreuzer, M., Bérard, J., & Hillmann, E. (2018). Long-term effects of castration, chronic intermittent social stress, provision of grass silage and their interactions on performance and meat and adipose tissue properties in growing-finishing pigs. *Meat Science*, 145, 40–50. <https://doi.org/10.1016/j.meatsci.2018.05.018>
- Horcada, A., Fernández-Cabanás, V. M., Polvillo, O., Botella, B., Dolores Cubiles, M., Pino, R., Narváez-Rivas, M., León-Camacho, M., & Acuña, R. R. (2013). Feasibility of use of fatty acid and triacylglycerol profiles for the authentication of commercial labelling in Iberian dry-cured sausages. *Talanta*, 117, 463–470. <https://doi.org/10.1016/j.talanta.2013.09.029>
- Horcada, A., Valera, M., Juárez, M., & Fernández-Cabanás, V. (2020). Authentication of Iberian pork official quality categories using a portable near infrared spectroscopy (NIRS) instrument. *Food Chemistry*, 318, 126471. <https://doi.org/10.1016/j.foodchem.2020.126471>
- Jaturasitha, S., Chaiwang, N., Kayan, A., & Kreuzer, M. (2016). Nutritional strategies to improve the lipid composition of meat, with emphasis on Thailand and Asia. *Meat Science*, 120, 157–166. <https://doi.org/10.1016/j.meatsci.2016.04.014>
- Juárez, M., Lam, S., Bohrer, B. M., Dugan, M. E. R., Vahmani, P., Aalhus, J., Juárez, A., López-Campos, O., Prieto, N., & Segura, J. (2021). Enhancing the Nutritional Value of Red Meat through Genetic and Feeding Strategies. *Foods*, 10(4), 872. <https://doi.org/10.3390/foods10040872>
- Larson-Meyer, D. E., Ingold, B. C., Fensterseifer, S. R., Austin, K. J., Wechsler, P. J., Hollis, B. W., Makowski, A. J., & Alexander, B. M. (2017). Sun exposure in pigs increases the

- vitamin D nutritional quality of pork. *PLOS ONE*, 12(11), e0187877.
<https://doi.org/10.1371/journal.pone.0187877>
- Lebret, B., Dourmad, J. Y., Mourot, J., Pollet, P. Y., & Gondret, F. (2014). Production performance, carcass composition, and adipose tissue traits of heavy pigs: Influence of breed and production system1. *Journal of Animal Science*, 92(8), 3543–3556.
<https://doi.org/10.2527/jas.2013-7398>
- Martins, J. M., Albuquerque, A., Neves, J. A., Freitas, A. B., Charneca, R., & Tirapicos, J. L. (2017). Influence of outdoor rearing and oleic acid supplementation on lipid characteristics of muscle and adipose tissues from obese Alentejano pigs. *Journal of Animal Physiology and Animal Nutrition*, 102(2), e578–e590. <https://doi.org/10.1111/jpn.12799>
- Martins, J., Neves, J., Freitas, A., & Tirapicos, J. (2015). Rearing system and oleic acid supplementation effect on carcass and lipid characteristics of two muscles from an obese pig breed. *Animal*, 9(10), 1721–1730. <https://doi.org/10.1017/s175173111500066x>
- N., O., N., C. R., Y., M., & M., O. E. (2018). Effects of a pasture-based pork production system on the expression of genes involved in lipid metabolism and meat quality characteristics. *International Journal of Nutrition and Metabolism*, 10(4), 23–30.
<https://doi.org/10.5897/ijnam2018.0232>
- Overholt, M. F., Lowell, J. E., Wilson, K. B., Matulis, R. J., Stein, H. H., Dilger, A. C., & Boler, D. D. (2016). Effects of feeding pelleted diets without or with distillers dried grains with solubles on fresh belly characteristics, fat quality, and commercial bacon slicing yields of finishing pigs. *Journal of Animal Science*, 94(5), 2198. <https://doi.org/10.2527/jas.2015-0203>
- Paredi, G., Mori, F., de Marino, M. G., Raboni, S., Marchi, L., Galati, S., Buschini, A., lo Fiego, D. P., & Mozzarelli, A. (2019). Is the protein profile of pig Longissimus dorsi affected by gender and diet? *Journal of Proteomics*, 206, 103437.
<https://doi.org/10.1016/j.jprot.2019.103437>
- Prévéraud, D. P., Desmarchelier, C., Rouffineau, F., Devillard, E., & Borel, P. (2015). A meta-analysis to assess the effect of the composition of dietary fat on α -tocopherol blood and tissue concentration in pigs. *Journal of Animal Science*, 93(3), 1177.
<https://doi.org/10.2527/jas.2014-8422>
- Qi, K., Men, X., Wu, J., & Xu, Z. (2019). Rearing pattern alters porcine myofiber type, fat deposition, associated microbial communities and functional capacity. *BMC Microbiology*, 19(1). <https://doi.org/10.1186/s12866-019-1556-x>
- Reig, M., Aristoy, M. C., & Toldrá, F. (2013). Variability in the contents of pork meat nutrients and how it may affect food composition databases. *Food Chemistry*, 140(3), 478–482.
<https://doi.org/10.1016/j.foodchem.2012.11.085>

Ribeiro, D. M., Mourato, M. P., & Almeida, A. M. (2019). Assessing mineral status in edible tissues of domestic and game animals: a review with a special emphasis in tropical regions. *Tropical Animal Health and Production*, 51(5), 1019–1032. <https://doi.org/10.1007/s11250-019-01848-8>

Sánchez Del Pulgar, J., Carrapiso, A., Reina, R., Biasioli, F., & García, C. (2013). Effect of IGF-II genotype and pig rearing system on the final characteristics of dry-cured Iberian hams. *Meat Science*, 95(3), 586–592. <https://doi.org/10.1016/j.meatsci.2013.05.044>

Shircliff, K. E., Callahan, Z. D., Wilmoth, T. A., Ohman, C. E., Johnson, R. C., & Wiegand, B. R. (2015). Growth performance and pork fat quality as measured by three methods of pigs fed 20% DDGS and slaughtered using a standard industry split marketing strategy. *Journal of Animal Science*, 93(10), 5083–5090. <https://doi.org/10.2527/jas.2015-9394>

Średnicka-Tober, D., Barański, M., Seal, C., Sanderson, R., Benbrook, C., Steinshamn, H., Gromadzka-Ostrowska, J., Rembiałkowska, E., Skwarło-Sońta, K., Eyre, M., Cozzi, G., Krogh Larsen, M., Jordon, T., Niggli, U., Sakowski, T., Calder, P. C., Burdge, G. C., Sotiraki, S., Stefanakis, A., . . . Leifert, C. (2016a). Composition differences between organic and conventional meat: a systematic literature review and meta-analysis. *British Journal of Nutrition*, 115(6), 994–1011. <https://doi.org/10.1017/s0007114515005073>

Średnicka-Tober, D., Barański, M., Seal, C., Sanderson, R., Benbrook, C., Steinshamn, H., Gromadzka-Ostrowska, J., Rembiałkowska, E., Skwarło-Sońta, K., Eyre, M., Cozzi, G., Krogh Larsen, M., Jordon, T., Niggli, U., Sakowski, T., Calder, P. C., Burdge, G. C., Sotiraki, S., Stefanakis, A., . . . Leifert, C. (2016b). Composition differences between organic and conventional meat: a systematic literature review and meta-analysis. *British Journal of Nutrition*, 115(6), 994–1011. <https://doi.org/10.1017/s0007114515005073>

Sheep and goats

Acciario, M., Decandia, M., Sitzia, M., Manca, C., Giovanetti, V., Cabiddu, A., Addis, M., Rassu, S. P. G., Molle, G., & Dimauro, C. (2020). Discriminant analysis as a tool to identify bovine and ovine meat produced from pasture or stall-fed animals. *Italian Journal of Animal Science*, 19(1), 1065–1070. <https://doi.org/10.1080/1828051x.2020.1816507>

Aguayo-Ulloa, L., Pascual-Alonso, M., Campo, M., Olleta, J., Villarroel, M., Pizarro, D., Miranda-de La Lama, G., & María, G. (2014). Effects of an enriched housing environment on sensory aspects and fatty-acid composition of the longissimus muscle of light-weight finished lambs. *Meat Science*, 97(4), 490–496. <https://doi.org/10.1016/j.meatsci.2014.02.017>

Álvarez, R., Meléndez-Martínez, A., Vicario, I., & Alcalde, M. (2014). Effect of pasture and concentrate diets on concentrations of carotenoids, vitamin A and vitamin E in plasma and adipose tissue of lambs. *Journal of Food Composition and Analysis*, 36(1–2), 59–65. <https://doi.org/10.1016/j.jfca.2014.08.001>

- Bernard, L., Fougère, H., Larsen, T., & Pires, J. (2020). Short communication: Diets supplemented with starch and corn oil, marine algae, or hydrogenated palm oil differently affect selected metabolite concentrations in cow and goat milk. *Journal of Dairy Science*, *103*(6), 5647–5653. <https://doi.org/10.3168/jds.2019-18008>
- Bravo-Lamas, L., Barron, L. J., Kramer, J. K., Etaio, I., & Aldai, N. (2016). Characterization of the fatty acid composition of lamb commercially available in northern Spain: Emphasis on the trans-18:1 and CLA content and profile. *Meat Science*, *117*, 108–116. <https://doi.org/10.1016/j.meatsci.2016.02.043>
- Cabrera, M., & Saadoun, A. (2014). An overview of the nutritional value of beef and lamb meat from South America. *Meat Science*, *98*(3), 435–444. <https://doi.org/10.1016/j.meatsci.2014.06.033>
- Cais-Sokolińska, D., Wójtowski, J., Pikul, J., Danków, R., Majcher, M., Teichert, J., & Bagnicka, E. (2015). Formation of volatile compounds in kefir made of goat and sheep milk with high polyunsaturated fatty acid content. *Journal of Dairy Science*, *98*(10), 6692–6705. <https://doi.org/10.3168/jds.2015-9441>
- Cardoso, D. B., Vêras, R. M. L., de Carvalho, F. F. R., Magalhães, A. L. R., de Vasconcelos, G. A., Maciel, M. I. S., Madruga, M. S., Urbano, S. A., & de Lima Silva, J. (2019). Physicochemical parameters, fatty acid profile, and sensory attributes of meat from lambs fed with cassava dregs in replacement of corn. *Tropical Animal Health and Production*, *51*(6), 1515–1521. <https://doi.org/10.1007/s11250-019-01840-2>
- Cherif, M., Valenti, B., Abidi, S., Luciano, G., Mattioli, S., Pauselli, M., Bouzarraa, I., Priolo, A., & ben Salem, H. (2018). Supplementation of *Nigella sativa* seeds to Barbarine lambs raised on low- or high-concentrate diets: Effects on meat fatty acid composition and oxidative stability. *Meat Science*, *139*, 134–141. <https://doi.org/10.1016/j.meatsci.2018.01.022>
- Chikwanha, O. C., Vahmani, P., Muchenje, V., Dugan, M. E., & Mapiye, C. (2018). Nutritional enhancement of sheep meat fatty acid profile for human health and wellbeing. *Food Research International*, *104*, 25–38. <https://doi.org/10.1016/j.foodres.2017.05.005>
- Cividini, A., Levart, A., Žgur, S., & Kompan, D. (2014). Fatty acid composition of lamb meat from the autochthonous Jezersko–Solčava breed reared in different production systems. *Meat Science*, *97*(4), 480–485. <https://doi.org/10.1016/j.meatsci.2013.12.012>
- DARWISH, W. S., IKENAKA, Y., MORSHDY, A. E., ELDESOKY, K. I., NAKAYAMA, S., MIZUKAWA, H., & ISHIZUKA, M. (2016). β -carotene and retinol contents in the meat of herbivorous ungulates with a special reference to their public health importance. *Journal of Veterinary Medical Science*, *78*(2), 351–354. <https://doi.org/10.1292/jvms.15-0287>
- Dong, H., Wang, S., Jia, Y., Ni, Y., Zhang, Y., Zhuang, S., Shen, X., & Zhao, R. (2013). Long-Term Effects of Subacute Ruminal Acidosis (SARA) on Milk Quality and Hepatic Gene Expression in Lactating Goats Fed a High-Concentrate Diet. *PLoS ONE*, *8*(12), e82850. <https://doi.org/10.1371/journal.pone.0082850>

- Erasmus, S. W., Muller, M., & Hoffman, L. C. (2017). Authentic sheep meat in the European Union: Factors influencing and validating its unique meat quality. *Journal of the Science of Food and Agriculture*, 97(7), 1979–1996. <https://doi.org/10.1002/jsfa.8180>
- Fonteles, N. L., Alves, S. P., Madruga, M. S., Queiroga, R. R., Andrade, A. P., Silva, D. S., Leal, A. P., Bessa, R. J., & Medeiros, A. N. (2018). Fatty acid composition of polar and neutral meat lipids of goats browsing in native pasture of Brazilian Semiarid. *Meat Science*, 139, 149–156. <https://doi.org/10.1016/j.meatsci.2018.01.021>
- Hajji, H., Prache, S., Andueza, D., Smeti, S., Mahouachi, M., & Atti, N. (2019). Reliability of visible reflectance spectroscopy in discriminating between pasture and stall-fed lambs from thin and fat-tailed sheep breeds in dry and hot environment. *Animal*, 13(11), 2669–2678. <https://doi.org/10.1017/s1751731119000909>
- Howes, N. L., Bekhit, A. E. D. A., Burritt, D. J., & Campbell, A. W. (2014). Opportunities and Implications of Pasture-Based Lamb Fattening to Enhance the Long-Chain Fatty Acid Composition in Meat. *Comprehensive Reviews in Food Science and Food Safety*, 14(1), 22–36. <https://doi.org/10.1111/1541-4337.12118>
- Huang, Y., Andueza, D., de Oliveira, L., Zawadzki, F., & Prache, S. (2015). Visible spectroscopy on carcass fat combined with chemometrics to distinguish pasture-fed, concentrate-fed and concentrate-finished pasture-fed lambs. *Meat Science*, 101, 5–12. <https://doi.org/10.1016/j.meatsci.2014.10.024>
- Jacob, R., & Pethick, D. (2014). Animal factors affecting the meat quality of Australian lamb meat. *Meat Science*, 96(2), 1120–1123. <https://doi.org/10.1016/j.meatsci.2013.10.039>
- Jittakhot, S. (2014). Fatty acid composition in adipose tissues of goats and sheep reared on either grazing or confinement regimen. *Journal of Applied Animal Science*, 7(3), 23–30. <https://www.semanticscholar.org/paper/Fatty-acid-composition-in-adipose-tissues-of-goats-Jittakhot-Homyog/61c1934826639f55ed08fa640c209ab004bcdcb1>
- Leticia, M., Paola, D., Jordi, O., Julio, O., & Sancho, B. (2017). Effects of sage distillation by-product (*Salvia lavandulifolia* Vahl.) dietary supplementation in light lambs fed on concentrates on meat shelf life and fatty acid composition. *Meat Science*, 134, 44–53. <https://doi.org/10.1016/j.meatsci.2017.07.007>
- López-Andrés, P., Luciano, G., Vasta, V., Gibson, T., Scerra, M., Biondi, L., Priolo, A., & Mueller-Harvey, I. (2014). Antioxidant effects of ryegrass phenolics in lamb liver and plasma. *Animal*, 8(1), 51–57. <https://doi.org/10.1017/s1751731113001821>
- Luciano, G., Biondi, L., Scerra, M., Serra, A., Mele, M., Lanza, M., & Priolo, A. (2013). The effect of the change from a herbage- to a concentrate-based diet on the oxidative stability of raw and cooked lamb meat. *Meat Science*, 95(2), 212–218. <https://doi.org/10.1016/j.meatsci.2013.05.015>

- Mateo, L., Ortuño, J., & Bañón, S. (2021). Ewe's diet during gestation and lactation affects ready-to-eat meat from light lambs. *Animal*, *15*(2), 100043. <https://doi.org/10.1016/j.animal.2020.100043>
- Ocak, S., Ogun, S., & Yilmaz, O. (2016). Dorper sheep utilizing feed resources efficiently: a Mediterranean case study. *Revista Brasileira de Zootecnia*, *45*(8), 489–498. <https://doi.org/10.1590/s1806-92902016000800010>
- Pellattiero, E., Cecchinato, A., Tagliapietra, F., Schiavon, S., & Bittante, G. (2015). Determination by GC×GC of Fatty Acid and Conjugated Linoleic Acid (CLA) Isomer Profiles in Six Selected Tissues of Lambs Fed on Pasture or on Indoor Diets with and without Rumen-Protected CLA. *Journal of Agricultural and Food Chemistry*, *63*(3), 963–974. <https://doi.org/10.1021/jf504956x>
- Ponnampalam, E. N., Butler, K. L., Jacob, R. H., Pethick, D. W., Ball, A. J., Hocking Edwards, J. E., Geesink, G., & Hopkins, D. L. (2014). Health beneficial long chain omega-3 fatty acid levels in Australian lamb managed under extensive finishing systems. *Meat Science*, *96*(2), 1104–1110. <https://doi.org/10.1016/j.meatsci.2013.04.007>
- Ponnampalam, E. N., Butler, K. L., Pearce, K. M., Mortimer, S. I., Pethick, D. W., Ball, A. J., & Hopkins, D. L. (2014). Sources of variation of health claimable long chain omega-3 fatty acids in meat from Australian lamb slaughtered at similar weights. *Meat Science*, *96*(2), 1095–1103. <https://doi.org/10.1016/j.meatsci.2012.11.039>
- Ponnampalam, E., Vahedi, V., Giri, K., Lewandowski, P., Jacobs, J., & Dunshea, F. (2019). Muscle Antioxidant Enzymes Activity and Gene Expression Are Altered by Diet-Induced Increase in Muscle Essential Fatty Acid (α -linolenic acid) Concentration in Sheep Used as a Model. *Nutrients*, *11*(4), 723. <https://doi.org/10.3390/nu11040723>
- Realini, C., Bianchi, G., Bentancur, O., & Garibotto, G. (2017). Effect of supplementation with linseed or a blend of aromatic spices and time on feed on fatty acid composition, meat quality and consumer liking of meat from lambs fed dehydrated alfalfa or corn. *Meat Science*, *127*, 21–29. <https://doi.org/10.1016/j.meatsci.2016.12.013>
- Średnicka-Tober, D., Barański, M., Seal, C. J., Sanderson, R., Benbrook, C., Steinshamn, H., Gromadzka-Ostrowska, J., Rembiałkowska, E., Skwarło-Sońta, K., Eyre, M., Cozzi, G., Larsen, M. K., Jordon, T., Niggli, U., Sakowski, T., Calder, P. C., Burdige, G. C., Sotiraki, S., Stefanakis, A., . . . Leifert, C. (2016). Higher PUFA and n-3 PUFA, conjugated linoleic acid, α -tocopherol and iron, but lower iodine and selenium concentrations in organic milk: a systematic literature review and meta- and redundancy analyses. *British Journal of Nutrition*, *115*(6), 1043–1060. <https://doi.org/10.1017/s0007114516000349>
- Średnicka-Tober, D., Barański, M., Seal, C., Sanderson, R., Benbrook, C., Steinshamn, H., Gromadzka-Ostrowska, J., Rembiałkowska, E., Skwarło-Sońta, K., Eyre, M., Cozzi, G., Krogh Larsen, M., Jordon, T., Niggli, U., Sakowski, T., Calder, P. C., Burdige, G. C., Sotiraki, S., Stefanakis, A., . . . Leifert, C. (2016). Composition differences between

- organic and conventional meat: a systematic literature review and meta-analysis. *British Journal of Nutrition*, 115(6), 994–1011. <https://doi.org/10.1017/s0007114515005073>
- Teng, F., Reis, M. G., Broadhurst, M., Lagutin, K., Samuelsson, L., Ma, Y., Stevens, D., & Day, L. (2020). Factors affecting levels of volatile 4-alkyl branched-chain fatty acids in sheep milk from 2 contrasting farming systems in New Zealand. *Journal of Dairy Science*, 103(3), 2419–2433. <https://doi.org/10.3168/jds.2019-17192>
- Toral, P., Monahan, F., Hervás, G., Frutos, P., & Moloney, A. (2018). Review: Modulating ruminal lipid metabolism to improve the fatty acid composition of meat and milk. Challenges and opportunities. *Animal*, 12, s272–s281. <https://doi.org/10.1017/s1751731118001994>
- Valdivielso, I., Bustamante, M. N., Buccioni, A., Franci, O., Ruiz De Gordo, J. C., de Renobales, M., & Barron, L. J. R. (2015). Commercial sheep flocks – fatty acid and fat-soluble antioxidant composition of milk and cheese related to changes in feeding management throughout lactation. *Journal of Dairy Research*, 82(3), 334–343. <https://doi.org/10.1017/s0022029915000369>
- van Leeuwen, K. A., Camin, F., Jerónimo, E., Vasta, V., Prenzler, P. D., Ryan, D., & Bessa, R. J. B. (2017). Dietary Effects on Stable Carbon Isotope Composition of Fatty Acids in Polar and Neutral Fractions of Intramuscular Fat of Lambs. *Journal of Agricultural and Food Chemistry*, 65(43), 9404–9411. <https://doi.org/10.1021/acs.jafc.7b02999>
- van Vliet, S., Provenza, F. D., & Kronberg, S. L. (2021). Health-Promoting Phytonutrients Are Higher in Grass-Fed Meat and Milk. *Frontiers in Sustainable Food Systems*, 4. <https://doi.org/10.3389/fsufs.2020.555426>
- Wang, B., Luo, Y., Wang, Y., Wang, D., Hou, Y., Yao, D., Tian, J., & Jin, Y. (2020). Rumen bacteria and meat fatty acid composition of Sunit sheep reared under different feeding regimens in China. *Journal of the Science of Food and Agriculture*, 101(3), 1100–1110. <https://doi.org/10.1002/jsfa.10720>
- Wang, J., Xu, Z., Zhang, H., Wang, Y., Liu, X., Wang, Q., Xue, J., Zhao, Y., & Yang, S. (2021). Meat differentiation between pasture-fed and concentrate-fed sheep/goats by liquid chromatography quadrupole time-of-flight mass spectrometry combined with metabolomic and lipidomic profiling. *Meat Science*, 173, 108374. <https://doi.org/10.1016/j.meatsci.2020.108374>
- Watkins, P. J., Frank, D., Singh, T. K., Young, O. A., & Warner, R. D. (2013). Sheepmeat Flavor and the Effect of Different Feeding Systems: A Review. *Journal of Agricultural and Food Chemistry*, 61(15), 3561–3579. <https://doi.org/10.1021/jf303768e>
- Webb, E., & Erasmus, L. (2014). The effect of production system and management practices on the quality of meat products from ruminant livestock. *South African Journal of Animal Science*, 43(3), 413. <https://doi.org/10.4314/sajas.v43i3.12>

Whitney, T., & Smith, S. (2015). Substituting redberry juniper for oat hay in lamb feedlot diets: Carcass characteristics, adipose tissue fatty acid composition, and sensory panel traits. *Meat Science*, *104*, 1–7. <https://doi.org/10.1016/j.meatsci.2015.01.010>

Zhang, H., Ao, C., Khas-Erdene, Song, L., & Zhang, X. (2015). Effects of different model diets on milk composition and expression of genes related to fatty acid synthesis in the mammary gland of lactating dairy goats. *Journal of Dairy Science*, *98*(7), 4619–4628. <https://doi.org/10.3168/jds.2013-7097>